



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

1910. "A problem in voting." Edinburgh, Proc. Math. Soc., 28.  
 "Classification of geometries with projective metric." Edinburgh, Proc. Math. Soc., 28, 25-41.  
 "Elementary considerations relating to the absolute." Edinburgh, Proc. Math. Soc., 28, 65-72.  
 "Note on the geometries in which straight lines are represented by circles." Edinburgh, Proc. Math. Soc., 28, 81-94.  
 "The early history of non-euclidean geometry." *Nature*, London, 84, 172.  
 "On the need of a non-euclidean bibliography." Rep. Brit. Assoc., 80 (Sheffield), 531-532.
1911. "The numerically greatest term of a binomial expansion." Edinburgh, Math. Notes, No. 7, 74-77.  
 "Concrete representations of non-euclidean geometry." In a Collection of Scientific Papers published by the University of St. Andrews to commemorate the 500th anniversary of its foundation.  
 "Bibliography of non-euclidean geometry, including the theory of parallels, the foundations of geometry, and space of  $n$  dimensions." London: Harrison, pp. 400.

## NOTES ON GREATEST COMMON DIVISOR AND LEAST COMMON MULTIPLE OF INTEGERS.

By BENJAMIN FRANKLIN YANNEY, Wooster University.

### I. PROPERTIES OF QUOTIENTS.

Let  $q_1, q_2, \dots, q_n$  be the respective quotients obtained by dividing the positive integers  $a_1, a_2, \dots, a_n$  by their greatest common divisor  $D$ ; and  $q'_1, q'_2, \dots, q'_n$  the respective quotients obtained by dividing  $L$ , the lowest common multiple of the integers  $a_1, a_2, \dots, a_n$ , by the integers, in turn. Hence

$$q_1 = a_1 \div D; q_2 = a_2 \div D; \dots; q_n = a_n \div D \dots (1);$$

$$q'_1 = L \div a_1; q'_2 = L \div a_2; \dots; q'_n = L \div a_n \dots (2).$$

The two sets of quotients possess the following properties:

1. Neither set of quotients has a factor common to all the numbers in the set. For, otherwise,  $D$  would not be the greatest common divisor nor  $L$  the least common multiple of the numbers  $a_1, a_2, \dots, a_n$ .

2.  $q_1 q'_1 = q_2 q'_2 = \dots = q_n q'_n = L \div D$ .

This result is obtained by multiplying, member by member, the cor-

responding equalities in (1) and (2). Furthermore, by suitably combining all the equalities in (1) and (2), we get

$$3. (q_1 q_2 \dots q_n) (q'_1 q'_2 \dots q'_n) = L^n \div D^n; \text{ and,}$$

$$4. (a_1 a_2 \dots a_n)^2 = L^n D^n (q_1 q_2 \dots q_n) \div (q'_1 q'_2 \dots q'_n).$$

From the second property stated above it is easily seen that

5.  $L \div D$  divided in order by either set of quotients gives in corresponding order the other set.

From the first and fifth properties, it follows that

6. The least common multiple of each set of quotients is  $L \div D$ .

## II. RELATION OF INTEGERS TO THEIR GREATEST COMMON DIVISOR AND LEAST COMMON MULTIPLE.

1. As is well known, the product of two positive integers,  $a_1$  and  $a_2$ , is equal to the product of their greatest common divisor  $D$  and least common multiple  $L$ :

$$a_1 a_2 = DL.$$

The general relation may be stated as follows:

$$DL^{n-1} \supseteq a_1 a_2 \dots a_n \supseteq D^{n-1} L \dots (A).$$

PROOF. From (2) of I, we get

$$L^n = (a_1 a_2 \dots a_n) (q'_1 q'_2 \dots q'_n).$$

But, as previously shown, the least common multiple of  $q'_1, q'_2, \dots, q'_n$  is  $L \div D$ . Hence,  $q'_1 q'_2 \dots q'_n \supseteq L \div D$ . Therefore,  $L^n \supseteq (a_1 a_2 \dots a_n) L \div D$ .

$$\therefore DL^{n-1} \supseteq a_1 a_2 \dots a_n.$$

Similarly, starting with (1) of I, it can be shown that

$$a_1 a_2 \dots a_n \supseteq D^{n-1} L.$$

2. Another form of the general case is as follows:

$$\prod_{kD_i} \prod_{kL_i^{k-1}} \supseteq (a_1 a_2 \dots a_n) \frac{(n-1)!}{(n-k)!(k-1)!} \supseteq \prod_{kD_i^{k-1}} \prod_{kL_i} \dots (B),$$

in which  ${}_k D_i$  and  ${}_k L_i$  represent, respectively, the greatest common divisors and least common multiples of the  $n$  integers taken  $k$  at a time,  $i$  thus being

equal to 1, 2, ...,  $\frac{n!}{(n-k)!k!}$ .

The proof of this relation follows easily from (A), which is true for each set of  $k$  integers selected from the entire set of  $n$  integers. There will evidently be, in all,  $\frac{n!}{(n-k)!k!}$  such relations. Multiplying together these relations, member by member, and noticing that in the middle member each of the  $n$  integers will occur  $\frac{(n-1)!}{(n-k)!(k-1)!}$  times, the desired result is obtained.

3. If in (B) we set  $k=2$ , we obtain

$$(a_1 a_2 \dots a_n)^{n-1} = \prod {}_2D_i \prod {}_2L_i, \quad i=1, 2, \dots, \frac{1}{2}n(n-1),$$

which is probably the most general equality relation existing between integers, and greatest common divisors and least common multiples.

If, finally,  $n=2$ , we arrive at the well-known relation

$$a_1 a_2 = DL.$$

—————→—————

## A POINT'S VISIT TO THE LINEAR CONTINUUM.

By H. W. REDDICK, Columbia University.

Once upon a time a point, XYZ, in three-dimensional space, decided to visit the set of points on a certain straight line. He had a sense of his superiority over his less fortunate fellow-beings who were compelled to lead a one-dimensional existence, but his motive for visiting them might be called philanthropic for he had a real desire to find out the relations existing between them and to enlighten them, if possible, concerning a higher existence.

As XYZ approached the line he was surprised to find that its residents were not living happily together; and he soon learned that they were capable of the same petty dissensions and jealousies which he had supposed were possible only in the enlightened set to which he belonged. He found that none of them was on speaking terms with his next-door neighbor—in fact, that he did not know who his next-door neighbor was. The society was divided up into political parties, religious denominations, and exclusive cliques, which spent most of their time quarrelling with one another, and did not seem to realize that, taken all together, they formed one Grand Continuum.